

Title: Generating random variates from truncated distributions

Author: Wolfgang Bziuk

Short Description:

For some simulations random variates have to be generated from a truncated distribution with given mean. This can be done utilizing the non-truncated distribution, but often only the mean of the non-truncated random variables is adjustable such that it has to be modified properly.

Keywords: simulation, random variates, truncated distribution

Example:

Consider the nonnegative random variable \tilde{X} with mean value $E(\tilde{X})$, which is described through its distribution function $F_{\tilde{X}}(x) = \int_{x'=0}^x \tilde{f}_{\tilde{X}}(x')dx'$ and density $f_{\tilde{X}}(x)$ for $0 \leq x \leq \infty$. In order to generate a truncated distribution $F_X(x)$, $a \leq x \leq b$, with mean $E(X)$ the following methods are commonly used.

Method1: Scaled truncation

If the inverse transform method can be applied to $F_{\tilde{X}}(x)$ an algorithm for generating X is as follows [1]:

- Generate $U \sim U[0,1]$.
- Let $V = F_{\tilde{X}}(a) + [F_{\tilde{X}}(b) - F_{\tilde{X}}(a)] \cdot U$.
- Return $X = F_{\tilde{X}}^{-1}(V)$.

The truncated distribution function is given by

$$F_X(x) = \begin{cases} 0 & \text{for } x < a \\ \frac{F_{\tilde{X}}(x) - F_{\tilde{X}}(a)}{F_{\tilde{X}}(b) - F_{\tilde{X}}(a)} & \text{for } a \leq x \leq b \\ 1 & \text{for } x > b \end{cases}$$

Figure 1 gives an example for the exponential distribution.

Method2: Cut-off truncation

Assume we have a method for generating variates from $F_{\tilde{X}}(x)$. The truncation is done as follows:

- Generate X from $F_{\tilde{X}}(x)$.
- If $X \leq a$ let $X = a$.
- If $X \geq b$ let $X = b$.

In this case the truncated distribution function is given by [2]




Figure 1: Exponential distribution $\tilde{F}_X(x)$ and the scaled truncated distribution $F_X(x)$ with mean value $E(X)$.




Figure 2: Exponential distribution $\tilde{F}_X(x)$ and the cut-off truncated distribution $F_X(x)$ with mean value $E(X)$.

$$F_X(x) = \begin{cases} 0 & \text{für } x < a \\ F_{\tilde{X}}(x) & \text{für } a \leq x < b \\ 1 & \text{für } x \geq b \end{cases}$$

Figure 2 gives an example for the truncated exponential distribution.

References:

- [1] G. S. Fishman; 'Concepts And Methods In Diskrete Event Digital Simulation'; Wiley & Sons, 1973.
- [2] A. M. Law; W. D. Kelton; 'Simulation Modelling & Analysis', McGraw Hill Inc, 1991.